

# BUCHSBAUM HOMOGENEOUS ALGEBRAS WITH MINIMAL MULTIPLICITY

SHIRO GOTO AND KEN-ICHI YOSHIDA

ABSTRACT. In this paper we first give a lower bound on multiplicities for Buchsbaum homogeneous  $k$ -algebras  $A$  in terms of the dimension  $d$ , the codimension  $c$ , the initial degree  $q$ , and the length of the local cohomology modules of  $A$ . Next, we introduce the notion of Buchsbaum  $k$ -algebras with minimal multiplicity of degree  $q$ , and give several characterizations for those rings. In particular, we will show that those algebras have linear free resolutions. Further, we will give many examples of those algebras.

## 1. INTRODUCTION

In this paper we are going to introduce a new class of non-Cohen–Macaulay Buchsbaum rings with linear resolution. Now let us explain our motivation.

In 1967, Abhyankar [2] proved that for a homogeneous integral domain  $A$  over an algebraically closed field  $k$ , the following inequality holds:

$$\text{embdim}(A) = \dim_k A_1 \leq e(A) + \dim A - 1,$$

where  $e(A)$  (resp.  $\text{embdim}(A)$ ) denotes the multiplicity (resp. the embedding dimension) of  $A$ . In 1970, Sally [13] proved that the same inequality holds for any Cohen–Macaulay local ring, and called the ring  $A$  which satisfies the equality a *Cohen–Macaulay local ring with maximal embedding dimension*. In 1982, the first author [5] pointed out that Sally’s result can be extended to the class of Buchsbaum local rings. Namely, for any Buchsbaum local ring  $A$ , the following inequality holds:

$$\text{embdim}(A) \leq e(A) + \dim A - 1 + I(A),$$

where  $I(A)$  denotes the Buchsbaum invariant ( $I$ -invariant) of  $A$ . A Buchsbaum local ring  $A$  which satisfies the equality is said to be a *Buchsbaum local ring with maximal embedding dimension*. This is equivalent to that  $A$  has 2-linear resolution.

On the other hand, the first author [6] in 1983 proved that

$$e(A) \geq 1 + \sum_{i=1}^{d-1} \binom{d-1}{i-1} l_A(H_{\mathfrak{m}}^i(A)),$$

and defined the notion of *Buchsbaum local rings with minimal multiplicity*. He also proved that those rings have 2-linear resolutions.

Recently, Terai and the second author [16] defined the notion of *Buchsbaum rings with minimal multiplicity with initial degree  $q$  for Stanley–Reisner rings*, and proved that those rings have  $q$ -linear resolutions. Their result indicates the existence of Buchsbaum homogeneous  $k$ -algebras with minimal multiplicity of higher initial degree.

The main purpose of this paper is to introduce the notion of Buchsbaum homogeneous  $k$ -algebra with minimal multiplicity of higher (initial) degree and characterize those rings. Moreover, we will give several examples of those  $k$ -algebras.

---

*Date:* February 2, 2008.

*2000 Mathematics Subject Classification.* Primary 13H10; Secondary 13H15.

*Key words and phrases.* Buchsbaum ring, minimal multiplicity, regularity, linear resolution, initial degree, Stanley–Reisner ring.

Let us explain the organization of our paper. In Section 2, we recall several definitions: Buchsbaum property, initial degree, regularity, and  $a$ -invariant. We also recall two important theorems for Buchsbaum homogeneous  $k$ -algebras: Hoa–Miyazaki theorem, Eisenbud–Goto theorem. These theorems will play key roles in our proof of the main theorem.

In section 3, we give a lower bound on multiplicities for Buchsbaum homogeneous  $k$ -algebras in terms of its dimension, codimension and initial degree. Namely, we prove:

**Theorem 3.2 (Lower bound on multiplicities of Buchsbaum  $k$ -algebras).** Let  $A$  be a homogeneous Buchsbaum  $k$ -algebra with  $d = \dim A \geq 1$ ,  $c = \operatorname{codim} A \geq 1$  and  $q = \operatorname{indeg} A \geq 2$ . Let  $\mathfrak{M}$  denote the unique graded maximal ideal of  $A$ . Let  $Q$  be a parameter ideal of  $A$  which is generated by a linear system of parameters. Then

- (1)  $\Sigma(Q) \subseteq \mathfrak{M}^{q-1} + Q$ .
- (2) The following inequality holds:

$$e(A) \geq \binom{c+q-2}{q-2} + \sum_{i=1}^{d-1} \binom{d-1}{i-1} l_A(H_{\mathfrak{M}}^i(A)).$$

Moreover, the equality holds in (1) (or (2)) if and only if  $[\Sigma(Q)]_n = A_n$  for all  $n \geq q-1$ .

The ring  $A$  is said to be a *Buchsbaum  $k$ -algebra with minimal multiplicity of degree  $q$*  if the equality holds. Note that this notion coincides with having minimal multiplicity defined by the first author (resp. by Terai and the second author) when  $q = 2$  (resp. for Stanley–Reisner rings). Moreover, those algebras are *not* Cohen–Macaulay except polynomial rings. In fact, we know that  $e(A) \geq \binom{c+q-1}{q-1}$  if  $A$  is Cohen–Macaulay; see, e.g., Proposition 3.1.

In Section 4, we prove the following theorem, which gives a characterization of Buchsbaum  $k$ -algebras with minimal multiplicity of degree  $q$ .

**Theorem 4.1 (Characterization of Buchsbaum  $k$ -algebras with minimal multiplicity of degree  $q$ ).** Under the same notation as in Theorem 3.2, the following conditions are equivalent:

- (1)  $A$  has minimal multiplicity of degree  $q$ .
- (2)  $a(A) = q - d - 2$ .
- (3)  $H_{\mathfrak{M}}^i(A) = [H_{\mathfrak{M}}^i(A)]_{q-1-i}$  for all  $i < d$  and  $[H_{\mathfrak{M}}^d(A)]_n = 0$  for all  $n \geq q - d - 1$ .
- (4)  $A$  has  $q$ -linear resolution with

$$\sum_{i=0}^{d-1} \binom{d}{i} l_A(H_{\mathfrak{M}}^i(A)) = \binom{\operatorname{reg} A + c - 1}{c - 1}.$$

- (5)  $\Sigma(Q) = \mathfrak{M}^{q-1} + Q$ , that is,  $[\Sigma(Q)]_n = A_n$  ( $n \geq q-1$ ).

When this is the case, we have

- (a)  $\operatorname{reg} A = q - 1$ .
- (b)  $\operatorname{Soc}(H_{\mathfrak{M}}^d(A)) = [H_{\mathfrak{M}}^d(A)]_{q-d-2}$ .

In particular, the above theorem yields that any Buchsbaum  $k$ -algebra with minimal multiplicity of degree  $q$  has  $q$ -linear resolution. Furthermore, we recognize that such a algebra attains an upper bound on the inequality in Hoa–Miyazaki theorem, Kamoi–Vogel theorem, respectively; see Remark 4.9.

We also give some examples of Buchsbaum  $k$ -algebras  $A$  having minimal multiplicity with depth  $A = 0$ . For example, let  $S$  be a polynomial ring over a field  $k$  with  $\mathfrak{N}$  the graded maximal ideal of  $S$ . If  $S/J$  is a homogeneous Cohen–Macaulay  $k$ -algebra with  $(q-1)$ -linear resolution, then  $A = S/\mathfrak{N}J$  is a Buchsbaum  $k$ -algebra with minimal multiplicity of degree  $q$  with depth  $A = 0$ .

In Section 5, we give several examples of Buchsbaum  $k$ -algebras with minimal multiplicity of degree  $q$ . We first show that any non-Cohen–Macaulay, Buchsbaum  $k$ -algebra with  $e(A) \leq$

2 has minimal multiplicity of degree at most 3. Note that a Buchsbaum  $k$ -algebra with  $e(A) = 2$  does *not* necessarily have minimal multiplicity in the sense of Goto [6].

On the other hand, one can find many examples of reduced Buchsbaum  $k$ -algebras with minimal multiplicity of higher degree among Stanley–Reisner rings; see also [16]. On the other hand, we have no examples of those algebras which are integral domains over an algebraically closed field.

## 2. PRELIMINARIES

We first briefly recall the definition of Buchsbaum rings, which was introduced by Stückrad and Vogel. Let  $(A, \mathfrak{m})$  be a Noetherian local ring with the maximal ideal  $\mathfrak{m}$ , and  $M$  a finitely generated  $A$ -module of  $\dim M = s$ .

**Definition 2.1** (see, e.g., [15]). The module  $M$  is *Buchsbaum* if it satisfies one of the following equivalent conditions:

- (1) The difference  $l_A(M/QM) - e_Q(M)$  is independent of the choice of a parameter ideal  $Q$  for  $M$ , which we denote by  $I_A(M)$  and we call the Buchsbaum invariant of  $M$ .
- (2) Any system of parameters for  $M$  forms a weak  $M$ -sequence: For any system  $a_1, \dots, a_s$  of parameters for  $M$  and for any integer  $1 \leq i \leq s$ , one has the equality

$$(a_1, \dots, a_s)M : a_i = (a_1, \dots, a_{i-1})M : \mathfrak{m}.$$

- (3) Any system of parameters for  $M$  forms a  $d$ -sequence for  $M$  in the sense of Huneke, that is, the equality

$$(a_1, \dots, a_{i-1})M : a_i a_j = (a_1, \dots, a_{i-1})M : a_j$$

holds for all integers  $1 \leq i \leq j \leq s$ .

We say that  $A$  is a *Buchsbaum ring* if  $A$  is a Buchsbaum module over itself. The module  $M$  is *Cohen–Macaulay* if it is Buchsbaum with  $I_A(M) = 0$ .

Now let  $M$  be a Buchsbaum  $A$ -module of  $\dim M = s$ . Then the local cohomology module  $H_{\mathfrak{m}}^i(M)$  is a finite-dimensional vector space over  $A/\mathfrak{m}$  for all  $i \neq s$ . In particular,  $\mathfrak{m}H_{\mathfrak{m}}^i(M) = 0$  for all  $i \neq s$  and  $M_{\mathfrak{p}}$  is a Cohen–Macaulay  $A_{\mathfrak{p}}$ -module for all prime  $\mathfrak{p} \in \text{Supp } M \setminus \{\mathfrak{m}\}$ . Furthermore, the following formula is known for Buchsbaum invariant:

$$I_A(M) = \sum_{i=0}^{s-1} \binom{s-1}{i} l_A(H_{\mathfrak{m}}^i(M)).$$

Moreover, if  $a_1, \dots, a_r \in \mathfrak{m}$  is part of a system of parameters for  $M$ , then the quotient module  $\overline{M} = M/(a_1, \dots, a_r)M$  is also a Buchsbaum  $A$ -module and  $I_A(M) = I_A(\overline{M})$ .

The main target in this paper are homogeneous Buchsbaum  $k$ -algebras with linear resolutions. In the following, let us recall several fundamental definitions of homogeneous  $k$ -algebras. Let  $k$  be an infinite field, and let  $S = k[X_1, \dots, X_v]$  be a homogeneous polynomial ring with  $n$  variables over  $k$  with  $\deg X_i = 1$ . Let  $A = S/I$  be a  $d$ -dimensional homogeneous  $k$ -algebra, where  $I$  is a graded ideal such that  $0 \neq I \subseteq (X_1, \dots, X_v)^2 S$ . Take a graded minimal free resolution over  $S$ :

$$0 \rightarrow \bigoplus_{j \in \mathbb{Z}} S(-j)^{\beta_{p,j}(A)} \xrightarrow{\varphi_p} \dots \xrightarrow{\varphi_2} \bigoplus_{j \in \mathbb{Z}} S(-j)^{\beta_{1,j}(A)} \xrightarrow{\varphi_1} S \rightarrow A \rightarrow 0,$$

where  $\beta_{i,j}(A)$  denotes the graded Betti number of  $A$  and  $S(-j)$  denotes the graded free  $S$ -module with grading given by  $[S(-j)]_n = S_{n-j}$ . Then the *initial degree* of  $A$  is defined by

$$\text{indeg } A = \min\{j \in \mathbb{Z} : \beta_{1,j}(A) \neq 0\}.$$

Let  $\text{indeg } I$  be the smallest degree of minimal system of generators of  $I$ . Note that  $\text{indeg } S/I = \text{indeg } I$  provided that  $0 \neq I \subseteq (X_1, \dots, X_v)^2 S$ . Similarly, the (Castelnuovo–Mumford) *regularity* is defined by

$$\text{reg } A = \max\{j - i \in \mathbb{Z} : \beta_{i,j}(A) \neq 0\}.$$

Then  $\text{reg } A \geq \text{indeg } A - 1$ , and  $A$  has *q-linear resolution* if the equality holds and  $\text{indeg } A = q$ . Note that we consider  $\text{indeg } A = 1$  and  $\text{reg } A = 0$  provided that  $A(= S)$  is a polynomial ring.

Put  $\mathfrak{M} = (X_1, \dots, X_v)A$ , the unique homogeneous maximal ideal of  $A$ . Let  $H_{\mathfrak{M}}^i(A)$  denote the  $i$ th local cohomology module of  $A$ . This module has a natural graded structure. We recall the *a-invariant* of  $A$ :

$$a(A) = \max\{n \in \mathbb{Z} : [H_{\mathfrak{M}}^d(A)]_n \neq 0\}.$$

One can compute the regularity of  $A$  as follows (see [3]):

$$\text{reg } A = \min\{n \in \mathbb{Z} : [H_{\mathfrak{M}}^i(A)]_j = 0 \text{ for all } i + j > n\}.$$

In particular,  $a(A) + d \leq \text{reg } A$  holds, and the equality holds if  $A$  is Cohen–Macaulay.

The homogeneous  $k$ -algebra  $A$  is *Buchsbaum* if the local ring  $A_{\mathfrak{M}}$  is Buchsbaum. Furthermore, we use  $I(A)$  in replace of  $I(A_{\mathfrak{M}})$ . As for regularity for Buchsbaum  $k$ -algebras, the following theorem is known:

**Theorem 2.2** (Hoa–Miyazaki [10]). *Suppose that  $A$  is a homogeneous Buchsbaum  $k$ -algebra with  $d = \dim A$ . Then  $\text{reg } A \leq a(A) + d + 1$ .*

*In particular,  $\text{reg } A = a(A) + d$ , or  $a(A) + d + 1$ .*

**Corollary 2.3.** *Under the same notation as in Theorem 2.2, we have  $a(A) \geq q - d - 2$ .*

*Proof.* By definition and Theorem 2.2, we have

$$q - 1 = \text{indeg } A - 1 \leq \text{reg } A \leq a(A) + d + 1.$$

Thus we get the required inequality.  $\square$

Now let  $A = S/I$  be a homogeneous Buchsbaum  $k$ -algebra, and let  $a_1, \dots, a_d \in A_1$  be a linear system of parameters of  $A$  and fix it. For an integer  $\ell \geq 1$  we set

$$Q := (a_1, \dots, a_d)A, \quad \Sigma(\underline{a}^\ell) := \sum_{i=1}^d (a_1^\ell, \dots, \widehat{a_i^\ell}, \dots, a_d^\ell) : a_i^\ell + (a_1^\ell, \dots, a_d^\ell).$$

In particular, we write  $\Sigma(Q) = \Sigma(\underline{a})$  for simplicity. Then  $Q$  is a minimal reduction of  $\mathfrak{M}$ , that is,  $\mathfrak{M}^{r+1} = Q\mathfrak{M}^r$  for some integer  $r \geq 0$ . The following result plays a key role in this paper.

**Theorem 2.4** (see [6, Theorem 4.1]). *Suppose that  $(A, \mathfrak{m})$  is a Buchsbaum local ring or a homogeneous  $k$ -algebra, and that  $Q$  is a minimal reduction of  $\mathfrak{m}$ . Then we have*

$$e(A) = e_Q(A) = l_A(A/\Sigma(Q)) + \sum_{i=1}^{d-1} \binom{d-1}{i-1} l_A(H_{\mathfrak{m}}^i(A)).$$

Suppose that  $A = S/I$  has  $q$ -linear resolution. Then one can take a minimal system of generators of  $I$  whose degree is equal to  $q$  and thus  $I \subseteq \mathfrak{N}^q$ , where  $\mathfrak{N} = (X_1, \dots, X_v)S$ . Moreover,  $A/Q$  also has  $q$ -linear resolution, that is,  $A/Q \cong k[Y_1, \dots, Y_c]/(Y_1, \dots, Y_c)^q$ , where  $c = \text{codim } A$ . In the case of Buchsbaum rings, the following criterion for having  $q$ -linear resolution is known:

**Theorem 2.5** (Eisenbud–Goto [3, Corollary 1.15]). *Suppose that  $A$  is a homogeneous Buchsbaum  $k$ -algebra with  $\dim A = d$ . Then the following conditions are equivalent:*

- (1)  $A$  has  $q$ -linear resolution.
- (2)  $H_{\mathfrak{M}}^i(A) = [H_{\mathfrak{M}}^i(A)]_{q-1-i}$  for all  $i < d$  and  $[H_{\mathfrak{M}}^d(A)]_n = 0$  for all  $n \geq q - d$ .
- (3)  $\mathfrak{M}^q = Q\mathfrak{M}^{q-1}$ .

### 3. LOWER BOUND ON MULTIPLICITIES

In [6], the first author proved an inequality

$$(3.1) \quad e(A) \geq 1 + \sum_{i=1}^{d-1} \binom{d-1}{i-1} l_A(H_{\mathfrak{m}}^i(A))$$

for any  $d$ -dimensional Buchsbaum local ring  $(A, \mathfrak{m})$  and called it a *Buchsbaum local ring with minimal multiplicity* if the equality holds. The main purpose of this section is to prove an improved version of the above inequality for homogeneous Buchsbaum  $k$ -algebras in terms of  $d = \dim A$ ,  $c = \operatorname{codim} A$  and  $q = \operatorname{indeg} A$  and to define the notion of *Buchsbaum homogeneous  $k$ -algebras with minimal multiplicity of degree  $q$* .

Throughout this section, we use the following notation, unless otherwise specified: Let  $S = k[X_1, \dots, X_v]$  is a polynomial ring and  $I$  is an ideal of  $S$  with  $q := \operatorname{indeg} I \geq 2$ , and let  $A = S/I$  be a homogeneous  $k$ -algebra with  $d := \dim A \geq 1$ ,  $c := \operatorname{codim} A \geq 1$  and  $\operatorname{indeg} A = q$ . Let  $\mathfrak{M}$  (resp.  $\mathfrak{N}$ ) denote the unique graded maximal ideal of  $A$  (resp.  $S$ ). Also, let  $Q$  be a parameter ideal of  $A$  which is generated by a linear system of parameters  $a_1, \dots, a_d$ .

Before stating our result, let us recall a lower bound on multiplicities for Cohen–Macaulay  $k$ -algebras. Under the above notation, we have

$$l_A(A/Q) \geq \dim_k k[Y_1, \dots, Y_c]/(Y_1, \dots, Y_c)^q = \binom{c+q-1}{q-1}$$

since  $A/Q$  is isomorphic to  $k[Y_1, \dots, Y_c]/L$ , where  $L$  is an ideal such that  $L \subseteq (Y_1, \dots, Y_c)^q$ . Further, the equality holds if and only if  $a(A/Q) = q-1$ . From this one can easily obtain the following (see also [3]).

**Proposition 3.1.** *Under the above notation, we also suppose that  $A$  is Cohen–Macaulay. Then*

- (1)  $e(A) \geq \binom{c+q-1}{q-1}$ .
- (2)  $a(A) = \operatorname{reg} A - d \geq q - d - 1$ .
- (3) *The following conditions are equivalent:*
  - (a)  $e(A) = \binom{c+q-1}{q-1}$ .
  - (b)  $a(A) = q - d - 1$ .
  - (c)  $A$  has  $q$ -linear resolution.

*When this is the case,  $A/Q \cong k[Y_1, \dots, Y_c]/(Y_1, \dots, Y_c)^q$ , where  $k[Y_1, \dots, Y_c]$  is a polynomial ring in  $c$  variables over  $k$ .*

The following theorem is the first main result in this paper, which generalizes the inequality in Eq.(3.1) and in Proposition 3.1.

**Theorem 3.2.** *Under the above notation, we also suppose that  $A$  is Buchsbaum. Then*

- (1)  $\Sigma(Q) \subseteq \mathfrak{M}^{q-1} + Q$ .
- (2) *The following inequality holds:*

$$e(A) \geq \binom{c+q-2}{q-2} + \sum_{i=1}^{d-1} \binom{d-1}{i-1} l_A(H_{\mathfrak{M}}^i(A)).$$

*Moreover, the equality holds in (1) (or (2)) if and only if  $[\Sigma(Q)]_n = A_n$  for all  $n \geq q-1$ .*

*Proof.* (1) It suffices to show that  $[\Sigma(Q)]_n = [Q]_n$  for all  $n \leq q-2$ . Fix  $i$  with  $1 \leq i \leq d$ . Set  $B = A/(a_1, \dots, \hat{a}_i, \dots, a_d)A$ . Then  $B$  is a 1-dimensional Buchsbaum  $k$ -algebra, and we can write as in the form  $B = k[Y_1, \dots, Y_{c+1}]/I_B$  and  $H_{\mathfrak{M}}^0(B) = U/I_B$  for some indeterminates  $Y_1, \dots, Y_{c+1}$  and some ideals  $I_B \subseteq U$  of  $k[Y_1, \dots, Y_{c+1}]$ . Then  $\operatorname{indeg} U \geq q-1$  since  $\mathfrak{M}U \subseteq I_B$

and  $\text{indeg } I_B \geq \text{indeg } I = q$ . Therefore if  $n \leq q - 2$  then

$$\left[ \frac{(a_1, \dots, \widehat{a_i}, \dots, a_d) : a_i}{(a_1, \dots, \widehat{a_i}, \dots, a_d)} \right]_n = [H_{\mathfrak{M}}^0(B)]_n = 0.$$

Namely,  $[(a_1, \dots, \widehat{a_i}, \dots, a_d) : a_i]_n = [(a_1, \dots, \widehat{a_i}, \dots, a_d)]_n \subseteq [Q]_n$  for every  $n$ . This implies that  $[\Sigma(Q)]_n \subseteq [Q]_n$  for all  $n \leq q - 2$ .

(2) By Theorem 2.4, it is enough to show  $l_A(A/\Sigma(Q)) \geq \binom{c+q-2}{q-2}$ . By (1), we have

$$l_A(A/\Sigma(Q)) \geq \sum_{n=0}^{q-2} \dim_k[A/\Sigma(Q)]_n = \sum_{n=0}^{q-2} \dim_k[A/Q]_n.$$

Since  $\text{indeg } A/Q \geq \text{indeg } A = q$ , we get

$$\sum_{n=0}^{q-2} \dim_k[A/Q]_n = \sum_{n=0}^{q-2} \binom{c+n-1}{n} = \binom{c+q-2}{q-2}.$$

It is easy to see the last assertion.  $\square$

*Remark 3.3.* Under the same notation as in Theorem 3.2, if, in addition,  $q > q' \geq 2$ , then

$$e(A) > \binom{c+q'-2}{q'-2} + \sum_{i=1}^{d-1} \binom{d-1}{i-1} l_A(H_{\mathfrak{M}}^i(A)).$$

Hence one can replace  $\text{indeg } A = q$  with  $\text{indeg } A \geq q$  in the above theorem.

By virtue of the above theorem, we can define the notion of Buchsbaum homogeneous  $k$ -algebras with minimal multiplicity, which generalizes the notion defined by Goto in [6].

**Definition 3.4 (Minimal multiplicity).** Under the same notation as in Theorem 3.2, the ring  $A$  is said to be a Buchsbaum homogeneous  $k$ -algebras with *minimal multiplicity of degree  $q$*  if the equality holds:

$$e(A) = \binom{c+q-2}{q-2} + \sum_{i=1}^{d-1} \binom{d-1}{i-1} l_A(H_{\mathfrak{M}}^i(A)).$$

We regard a polynomial ring as a Buchsbaum (Cohen–Macaulay)  $k$ -algebra with minimal multiplicity of degree 1.

For Buchsbaum local rings, we can prove a similar inequality as that in the above theorem. Namely, we have: Let  $(A, \mathfrak{m}, k)$  be a  $d$ -dimensional Buchsbaum local ring with  $c = \text{codim } A$  and suppose that the  $\mathfrak{m}$ -adic completion  $\widehat{A} = R/I$ , where  $(R, \mathfrak{n})$  is a complete regular local ring and  $I$  is an ideal of  $R$  such that  $I \subseteq \mathfrak{n}^q$ . Also, let  $Q$  be a minimal reduction of  $\mathfrak{m}$ . Then the same assertions in Theorem 3.2 hold; see the proof for [8]. Note that this yields [11, Theorem 2.2] and [6, Theorem 4.1] when  $q = 2$ .

However we do *not* know whether the associated graded ring  $\text{gr}_{\mathfrak{m}}(A) := \bigoplus_{n \geq 0} \mathfrak{m}^n / \mathfrak{m}^{n+1}$  of a Buchsbaum local ring with minimal multiplicity of degree  $q$  also has the same property (especially Buchsbaum property) except the case of  $q = 2$  yet. Namely, we do not have a complete answer to the following question below; See also [5]. So we do *not* introduce the notion of *Buchsbaum local ring with minimal multiplicity of degree  $q$*  here.

**Question 3.5.** Let  $(A, \mathfrak{m}, k)$  be a  $d$ -dimensional Buchsbaum local ring with  $\text{codim } A = c$ . Suppose that  $\widehat{A} = R/I$  such that  $I \subseteq \mathfrak{n}^q$ , where  $(R, \mathfrak{n})$  is a complete regular local ring and  $I$  is an ideal of  $R$ .

If  $\Sigma(Q) = \mathfrak{m}^{q-1} + Q$  (or,  $\mathfrak{m}^q = Q\mathfrak{m}^{q-1}$ ), then is the associated graded ring  $\text{gr}_{\mathfrak{m}}(A)$  of  $A$  Buchsbaum?

The inequality in Theorem 3.2 does *not* necessarily hold in general for non-Buchsbaum rings.

**Example 3.6.** Let  $k$  be any field, and let  $q, d \geq 2$  be integers.

Put  $A = k[X_0, X_1, \dots, X_d]/(X_0(X_0, \dots, X_d)^{q-1})$ . Then

- (1)  $A$  is a homogeneous  $k$ -algebra with  $\dim A = d$ ,  $\text{codim } A = 1$  and  $\text{indeg } A = q$  with the unique maximal ideal  $\mathfrak{m} = (x_0, \dots, x_d)A$ .
- (2)  $H_{\mathfrak{m}}^0(A) = x_0A$  and  $A/H_{\mathfrak{m}}^0(A) \cong k[X_1, \dots, X_d]$ . In particular,  $e(A) = 1$ .
- (3)  $A$  is Buchsbaum if and only if  $q = 2$ .
- (4)  $e(A) \geq \binom{c+q-2}{q-2} + \sum_{i=1}^{d-1} \binom{d-1}{i-1} l_A(H_{\mathfrak{m}}^i(A)) = q-1$  if and only if  $q = 2$ .

#### 4. CHARACTERIZATION

In this section, we give several equivalent conditions of Buchsbaum homogeneous  $k$ -algebras with minimal multiplicity of degree  $q$ . In particular, we show that those rings have  $q$ -linear resolutions and satisfy several boundary conditions; e.g., Hoa–Miyazaki theorem and Kamoi–Vogel theorem.

In this section, we use the same notation as in the previous section, unless otherwise specified. The following theorem is the second main result in this paper.

**Theorem 4.1.** *Let  $A$  be a homogeneous Buchsbaum  $k$ -algebra with  $d = \dim A \geq 1$ ,  $c = \text{codim } A \geq 1$  and  $q = \text{indeg } A \geq 2$ . Let  $Q$  be a parameter ideal of  $A$  which is generated by a linear system of parameters. Then the following conditions are equivalent:*

- (1)  $A$  has minimal multiplicity of degree  $q$ , that is,

$$e(A) = \binom{c+q-2}{q-2} + \sum_{i=1}^{d-1} \binom{d-1}{i-1} l_A(H_{\mathfrak{m}}^i(A)).$$

- (2)  $a(A) = q - d - 2$ .
- (3)  $H_{\mathfrak{m}}^i(A) = [H_{\mathfrak{m}}^i(A)]_{q-1-i}$  for all  $i < d$  and  $[H_{\mathfrak{m}}^d(A)]_n = 0$  for all  $n \geq q - d - 1$ .
- (4)  $A$  has  $q$ -linear resolution with

$$\sum_{i=0}^{d-1} \binom{d}{i} l_A(H_{\mathfrak{m}}^i(A)) = \binom{\text{reg } A + c - 1}{c - 1}.$$

- (5)  $\Sigma(Q) = \mathfrak{m}^{q-1} + Q$ , that is,  $[\Sigma(Q)]_n = A_n$  for all  $n \geq q - 1$ .

When this is the case, we have

- (a)  $\text{reg } A = q - 1$ .
- (b)  $\text{Soc}(H_{\mathfrak{m}}^d(A)) = [H_{\mathfrak{m}}^d(A)]_{q-d-2}$ .

In the following, let us prove Theorem 4.1. Before proving the theorem, let us prove the following lemma.

**Lemma 4.2.** *Suppose that  $A$  is a Buchsbaum  $k$ -algebra with  $q$ -linear resolution. Then  $[A/\Sigma(Q)]_{q-1} \cong [H_{\mathfrak{m}}^d(A)]_{q-d-1}$ .*

*Proof.* Now consider the following direct system:

$$(A/\Sigma(\underline{a})) (d) \longrightarrow (A/\Sigma(\underline{a}^2)) (2d) \longrightarrow \cdots \longrightarrow (A/\Sigma(\underline{a}^\ell)) (\ell d) \longrightarrow \cdots,$$

where the natural map  $(A/\Sigma(\underline{a}^k)) (kd) \longrightarrow (A/\Sigma(\underline{a}^\ell)) (\ell d)$  is given by a multiplication of  $(a_1 \cdots a_d)^{\ell-k}$  for  $1 \leq k < \ell$ . Then  $\varphi_\ell: (A/\Sigma(\underline{a}^\ell)) (\ell d) \longrightarrow H_{\mathfrak{m}}^d(A)$  is injective and  $\varinjlim (A/\Sigma(\underline{a}^\ell)) (\ell d) = H_{\mathfrak{m}}^d(A)$  since  $A$  is Buchsbaum; see [5] or [18, Lemma 1.3]. The lemma follows from the following claim.

**Claim.**  $A_{q-1+(\ell-1)d} \subseteq (a_1 \cdots a_d)^{(\ell-1)} \mathfrak{m}^{q-1} + \Sigma(\underline{a}^\ell)$  for every  $\ell \geq 1$ .

In fact, the claim follows from

$$\mathfrak{m}^{q-1+(\ell-1)d} = Q^{(\ell-1)d} \mathfrak{m}^{q-1} \subseteq (a_1^\ell, \dots, a_d^\ell) \mathfrak{m}^{q-1+(\ell-1)d-\ell} + (a_1 \cdots a_d)^{\ell-1} \mathfrak{m}^{q-1}.$$

□

*Proof of Theorem 4.1.* We have already seen  $(1) \iff (5)$  in the proof of Theorem 3.2. So it suffices to show  $(5) \implies (4) \implies (3) \implies (2) \implies (5)$ . Now we put  $h^i(A) = l_A(H_{\mathfrak{M}}^i(A))$  for all  $i < d$ .

$(5) \implies (4)$  : The assumption implies that  $\mathfrak{M}^q \subseteq \mathfrak{M}\Sigma(Q) \subseteq Q$ . This means that  $\mathfrak{M}^q = Q \cap \mathfrak{M}^q = Q\mathfrak{M}^{q-1}$  since  $A$  is homogeneous. Thus  $A$  has  $q$ -linear resolution by Eisenbud–Goto theorem (Theorem 2.5). In particular,  $\text{reg } A = q - 1$ , and we get

$$(4.1) \quad e(A) = e_Q(A) = l_A(A/Q) - I(A) = \binom{c+q-1}{q-1} - \sum_{i=0}^{d-1} \binom{d-1}{i} h^i(A).$$

On the other hand, by  $(5) \implies (1)$  we have

$$(4.2) \quad e(A) = \binom{c+q-2}{q-2} + \sum_{i=1}^{d-1} \binom{d-1}{i-1} h^i(A).$$

Comparing two equalities, we get

$$(4.3) \quad \sum_{i=0}^{d-1} \binom{d}{i} h^i(A) = \binom{c+q-2}{c-1} = \binom{\text{reg } A + c - 1}{c-1}.$$

$(4) \implies (3)$  : By a similar argument as above, we obtain that  $A$  has minimal multiplicity of degree  $q$ . In particular,  $[\Sigma(Q)]_{q-1} = A_{q-1}$ . This implies that  $[H_{\mathfrak{M}}^d(A)]_{q-1-d} = 0$  by Lemma 4.2 since  $A$  has  $q$ -linear resolution. On the other hand,  $H_{\mathfrak{M}}^i(A) = [H_{\mathfrak{M}}^i(A)]_{q-1-i}$  for all  $i < d$  and  $[H_{\mathfrak{M}}^d(A)]_n = 0$  for all  $n \geq q - d$  by Eisenbud–Goto theorem. Hence we get (3).

$(3) \implies (2)$  : By the assumption we have  $a(A) \leq q - d - 2$ . However the converse is always true; see Corollary 2.3.

$(2) \implies (5)$  : Assume that  $a(A) = q - d - 2$ . Then for all  $n \geq q - 1$ , we have

$$[A/\Sigma(Q)]_n \xrightarrow{\varphi_1} [H_{\mathfrak{M}}^d(A)]_{n-d} = 0.$$

Namely,  $[\Sigma(Q)]_n = A_n$  for all  $n \geq q - 1$ , as required.

We now prove that  $\text{Soc}(H_{\mathfrak{M}}^d(A)) = [H_{\mathfrak{M}}^d(A)]_{q-d-2}$  if  $a(A) = q - d - 2$ . It is enough to show that  $[\text{Soc}(H_{\mathfrak{M}}^d(A))]_n = 0$  for all  $n \leq q - d - 3$ . Consider the following direct system described as above:

$$[A/\Sigma(Q)]_{n+d} \xrightarrow{a_1 \cdots a_d} [A/\Sigma(\underline{a}^2)]_{n+2d} \xrightarrow{a_1 \cdots a_d} \cdots \longrightarrow [H_{\mathfrak{M}}^d(A)]_n.$$

By [18, Proposition 3.8], we have

$$\text{Soc}(H_{\mathfrak{M}}^d(A)) \subseteq \text{Hom}_A(A/Q, H_{\mathfrak{M}}^d(A)) = \varphi_2 \left( \frac{\sum_{i=1}^d a_1 \cdots \widehat{a_i} \cdots a_d \Sigma(Q) + \Sigma(\underline{a}^2)}{\Sigma(\underline{a}^2)} \right).$$

Since  $\Sigma(Q) = \mathfrak{M}^{q-1} + Q$ , if  $j \leq q + d - 3$ , then

$$\begin{aligned} \left[ \sum_{i=1}^d a_1 \cdots \widehat{a_i} \cdots a_d \Sigma(Q) + \Sigma(\underline{a}^2) \right]_j &= \sum_{i=1}^d a_1 \cdots \widehat{a_i} \cdots a_d [Q]_{j-d+1} + [\Sigma(\underline{a}^2)]_j \\ &\subseteq a_1 \cdots a_d A_{j-d} + [\Sigma(\underline{a}^2)]_j. \end{aligned}$$

Hence  $[\text{Soc}(H_{\mathfrak{M}}^d(A))]_n \subseteq [\text{Im } \varphi_1]_n$ , that is,  $[\text{Soc}(H_{\mathfrak{M}}^d(A))]_n = \left[ \frac{\Sigma(Q) : \mathfrak{M}}{\Sigma(Q)} \right]_{n+d}$  for all  $n \leq q - d - 3$ . It is enough to show the following claim.

**Claim.**  $[\Sigma(Q) : \mathfrak{M}]_j \subseteq [\Sigma(Q)]_j$  for all  $j \leq q - 3$ .

Fix  $j \leq q - 3$ . For any element  $\xi \in [\Sigma(Q) : \mathfrak{M}]_j$ ,  $\mathfrak{M}(\mathfrak{M}\xi) \subseteq \mathfrak{M}\Sigma(Q) \subseteq Q$ . Then for any element  $a \in A_1$ ,  $a\xi \in [\mathfrak{M}\xi]_{j+1} \subseteq [Q : \mathfrak{M}]_{j+1} = [Q]_{j+1}$  because  $\text{Soc}(A/Q)$  is concentrated in degree  $q - 1$ . Hence  $\mathfrak{M}\xi \subseteq Q$ . That is,  $\xi \in [Q : \mathfrak{M}]_j = [Q]_j \subseteq [\Sigma(Q)]_j$ , as required.

We have finished the proof of Theorem 4.1.  $\square$

The first author [7] determined the Hilbert series of Buchsbaum  $k$ -algebras with  $q$ -linear resolution. Thus we can also describe that of Buchsbaum  $k$ -algebras with minimal multiplicity.

**Corollary 4.3.** *Suppose that  $A$  is a Buchsbaum homogeneous  $k$ -algebras with minimal multiplicity of degree  $q$ . Put  $d = \dim A \geq 2$  and  $c = \operatorname{codim} A \geq 1$ . Then The Hilbert series  $F(A, t)$  of  $A$  is given as follows:*

$$F(A, t) = \frac{1}{(1-t)^d} \sum_{i=0}^{q-1} \binom{c+i-1}{i} t^i + \frac{1}{(1-t)^d} \sum_{i=q}^{q+d-1} (-1)^{i+q-1} \left\{ \sum_{j=0}^{q+d-1-i} \binom{d}{i+j-q+1} h^j(A) \right\} t^i.$$

Roughly speaking, a Buchsbaum homogeneous  $k$ -algebra with minimal multiplicity is *not* Cohen–Macaulay.

**Corollary 4.4.** *Suppose that  $A$  is a Buchsbaum homogeneous  $k$ -algebras with minimal multiplicity of degree  $q$ . Then  $A$  is Cohen–Macaulay if and only if it is isomorphic to a polynomial ring and  $q = 1$ .*

*Proof.* It follows from Proposition 3.1(2) and Theorem 4.1(2).  $\square$

In general,  $A/aA$  does *not* necessarily have minimal multiplicity even if so is  $A$  and  $a \in A_1$  is a non-zero-divisor.

**Corollary 4.5.** *Suppose that  $A$  is a  $d$ -dimensional Buchsbaum homogeneous  $k$ -algebras with minimal multiplicity of degree  $q$ . Let  $a \in A_1$  be a non-zero-divisor of  $A$  and set  $\bar{A} = A/aA$ . Then*

- (1)  $\bar{A}$  is a Buchsbaum  $k$ -algebra with  $q$ -linear resolution.
- (2)  $\bar{A}$  has minimal multiplicity if and only if  $H_{\mathfrak{M}}^{d-1}(A) = 0$ .

*Proof.* In the following, we put  $h^i(A) = l_A(H_{\mathfrak{M}}^i(A))$  for each  $i \leq d-1$ .

(1) The assumption implies that  $A$  has  $q$ -linear resolution by Theorem 4.1 (1)  $\implies$  (4). Hence  $\bar{A}$  also has  $q$ -linear resolution.

(2) We know that  $h^i(\bar{A}) = h^i(A) + h^{i+1}(A)$  for each  $i \leq d-2$  since  $A$  is Buchsbaum and  $a \in \mathfrak{M}$  is a non-zero-divisor. Hence

$$\sum_{i=1}^{d-2} \binom{d-2}{i-1} h^i(\bar{A}) = \sum_{i=1}^{d-1} \binom{d-1}{i-1} h^i(A) - h^{d-1}(A).$$

The assertion easily follows from this.  $\square$

Let us consider the case where  $A/H_{\mathfrak{M}}^0(A)$  is Cohen–Macaulay. Then one can obtain many examples of Buchsbaum homogeneous  $k$ -algebras with minimal multiplicity of degree  $q$ .

**Corollary 4.6.** *Under the same notation as in Theorem 4.1, if, in addition,  $A/H_{\mathfrak{M}}^0(A)$  is Cohen–Macaulay, then the following conditions are equivalent:*

- (1)  $A$  has minimal multiplicity of degree  $q$ , that is,  $e(A) = \binom{c+q-2}{q-2}$ .
- (2)  $A$  has  $q$ -linear resolution and  $l_A(H_{\mathfrak{M}}^0(A)) = \binom{c+q-2}{q-1}$ .
- (3)  $H_{\mathfrak{M}}^0(A) = [H_{\mathfrak{M}}^0(A)]_{q-1}$ ,  $H_{\mathfrak{M}}^i(A) = 0$  for all  $1 \leq i \leq d-1$  and  $[H_{\mathfrak{M}}^d(A)]_n = 0$  for all  $n \geq q-d-1$ .
- (4)  $A/H_{\mathfrak{M}}^0(A)$  has  $(q-1)$ -linear resolution.

*Proof.* It follows from Theorem 4.1 and Proposition 3.1.  $\square$

**Remark 4.7.** The condition (4) does *not* imply that  $A$  has minimal multiplicity (we need to assume  $\operatorname{indeg} A = q$ ); see also Example 5.6.

**Example 4.8.** If  $S/U$  is a Cohen–Macaulay  $k$ -algebra with  $(q - 1)$ -linear resolution, then  $A = S/\mathfrak{N}U$  is a Buchsbaum  $k$ -algebra with minimal multiplicity of degree  $q$  with  $l_A(H_{\mathfrak{N}}^0(A)) = \mu_S(U)$ , where  $\mu_S(U)$  denotes the minimal number of system of generators of  $U$ . For example,

- (1)  $A = S/f\mathfrak{N}$  is a Buchsbaum  $k$ -algebra with minimal multiplicity of degree  $q$  provided that  $f \in \mathfrak{N}^{q-1} \setminus \mathfrak{N}^q$ .
- (2) For given integers  $c \geq 1$ ,  $q \geq 2$ , if we put

$$A = k[X_1, \dots, X_c, Y_1, \dots, Y_d]/(X_1, \dots, X_c, Y_1, \dots, Y_d)(X_1, \dots, X_c)^{q-1},$$

then  $A$  is a Buchsbaum  $k$ -algebra with minimal multiplicity of degree  $q$  such that  $\dim A = d$ ,  $\operatorname{codim} A = c$  and  $A/H_{\mathfrak{N}}^0(A)$  is Cohen–Macaulay.

*Remark 4.9.* In [12], Kamoi and Vogel proved that the following inequality:

$$\sum_{i=0}^{d-1} \binom{d}{i} h^i(A) \leq \binom{\operatorname{reg} A + c - 1}{c - 1}$$

for any homogeneous Buchsbaum  $k$ -algebra  $A$ . Our theorem 4.1(3) shows that any Buchsbaum ring with minimal multiplicity attains the upper bound in this inequality. We do *not* know whether the above equality implies that  $A$  has linear resolution (and thus it has minimal multiplicity in our sense).

Furthermore, those rings satisfy “ $\operatorname{reg} A = a(A) + d + 1$ ” because  $\operatorname{reg} A = q - 1$  and  $a(A) = q - d - 2$ . See Hoa–Miyazaki theorem.

*Remark 4.10.* For a Buchsbaum  $k$ -algebra  $A$ , it has maximal embedding dimension if and only if it has 2-linear resolution.

It is known that any Buchsbaum ring with minimal multiplicity has 2-linear resolution; see [6]. Our theorem 4.1 (1)  $\iff$  (4) generalizes this fact.

## 5. EXAMPLES

In this section, we first show that all non-Cohen–Macaulay, Buchsbaum  $k$ -algebras with  $e(A) \leq 2$  have minimal multiplicities of degree at most 3. Next, we give several examples of Buchsbaum homogeneous  $k$ -algebras with minimal multiplicity of higher degree using the theory of Stanley–Reisner rings; see [16].

Throughout this section, let  $k$  be a field and  $S = k[X_1, \dots, X_v]$  a polynomial ring over  $k$  with the unique graded maximal ideal  $\mathfrak{N}$ .

**5.1. Buchsbaum rings with small multiplicities.** In [4], the first author classified non-Cohen–Macaulay, Buchsbaum rings  $(A, \mathfrak{m})$  with  $e(A) = 2$  and proved that such a ring has minimal multiplicity in the sense of Goto [6] such that  $h^i(A) : l_A(H_{\mathfrak{m}}^i(A)) = 0$  for all  $i \neq 1, d$  and  $h^1(A) = 1$  when  $d \geq 2$  and  $\operatorname{depth} A > 0$ ; see [4, Theorem 1.1] and also [6, Section 4] for details. But, when  $\operatorname{depth} A = 0$ , there exist a non-Cohen–Macaulay, Buchsbaum ring with  $e(A) = 2$  which does not have minimal multiplicity in the sense of Goto [6]. Now let us show that such a ring has minimal multiplicity in our sense. Before doing it, we prove the following lemma.

**Lemma 5.1.** *Suppose that  $A = S/I$  is a homogeneous non-Cohen–Macaulay, Buchsbaum  $k$ -algebra. Put  $d = \dim A \geq 2$ ,  $c = \operatorname{codim} A \geq 1$ ,  $q = \operatorname{indeg} A \geq 2$  and  $e = e(A)$ . Then*

- (1)  $q \leq e + 1$ .
- (2) *The following conditions are equivalent:*
  - (a)  $q = e + 1$ .
  - (b) *There exists an element  $f \in \mathfrak{N}^{q-1} \setminus \mathfrak{N}^q$  such that  $I = f\mathfrak{N}$ .*
  - (c)  *$A$  has minimal multiplicity of degree  $e + 1$ .*

*When this is the case,  $c = 1$  and  $h^i(A) = 0$  for all  $i = 1, \dots, d - 1$ .*

*Proof.* (1) Suppose that  $q \geq e + 2$ . By Theorem 3.2 we have

$$e \geq \binom{c+q-2}{q-2} \geq \binom{c+e}{e} \geq \binom{e+1}{e} = e + 1.$$

This is a contradiction. Hence  $q \leq e + 1$ .

(2) It is enough to show (a)  $\implies$  (b). We may assume that  $e \geq 2$ . Suppose (a). By Theorem 3.2 again, we have

$$e \geq \binom{c+e-1}{e-1} + \sum_{i=1}^{d-1} \binom{d-1}{i-1} h^i(A) \geq \binom{e}{e-1} + \sum_{i=1}^{d-1} \binom{d-1}{i-1} h^i(A) \geq e.$$

This implies that  $c = 1$ ,  $h^i(A) = 0$  for all  $i \neq 0, d$  and that  $A$  has minimal multiplicity of degree  $q = e + 1$ . Take an ideal  $U$  such that  $H_{\mathfrak{M}}^0(A) = U/I$ . Then  $\text{indeg } U \geq 2$  since  $\mathfrak{N}U \subseteq I$ . Hence  $\text{codim } A/H_{\mathfrak{M}}^0(A) = 1$  and thus  $A/H_{\mathfrak{M}}^0(A)$  is a hypersurface. So one can take an element  $f \in \mathfrak{N}$  such that  $U = fS$ . Since  $f\mathfrak{N} \subseteq I \subseteq fS$  and  $e(A/H_{\mathfrak{M}}^0(A)) = e = q - 1$ , we obtain (b).  $\square$

*Remark 5.2.* Most part of the above lemma is probably known. In fact, Hoa–Miyazaki theorem implies that  $q - 1 \leq \text{reg } A \leq a(A) + d + 1 \leq e(A)$  since  $A$  is Buchsbaum.

**Proposition 5.3.** *Suppose  $d \geq 2$ . Then any homogeneous non-Cohen–Macaulay, Buchsbaum  $k$ -algebra  $A$  with  $e(A) \leq 2$  has minimal multiplicity whose degree is at most 3. To be precise, we have:*

- (1) *Suppose that  $e(A) = 1$ . Then  $A$  has minimal multiplicity of degree 2 with  $\text{depth } A = 0$ . Moreover,  $A/H_{\mathfrak{M}}^0(A)$  is isomorphic to a polynomial ring over  $k$ .*
- (2) *Suppose that  $e(A) = 2$ . Then  $A$  has minimal multiplicity of degree  $q$  for some integer  $q \geq 2$ , and one of the following three cases occurs:*
  - (a)  $q = 2$  and  $\text{depth } A > 0$ . Then  $c = d$  and  $h^1(A) = 1$ .
  - (b)  $q = 2$  and  $\text{depth } A = 0$ . Then  $h^1(A) = 1$ .
  - (c)  $q = 3$  and  $\text{depth } A = 0$ . Then  $c = 1$  and  $A/H_{\mathfrak{M}}^0(A)$  is Cohen–Macaulay.

*Proof.* When  $\text{depth } A > 0$ , the assertion follows from [4, 6]. So we may suppose that  $\text{depth } A = 0$  and that  $q \geq 2$ .

(1) The assertion is clear. See also Theorem 3.2.

(2) By Lemma 5.1, we have  $q \leq 3$ .

**Case 1.** when  $h^1(A) \geq 1$ :

We obtain that  $q = 2$  by Lemma 5.1. Then  $A$  has minimal multiplicity and  $h^1(A) = 1$  since  $e(A) = 2 = 1 + h^1(A) \geq 2$ .

**Case 2.** when  $h^1(A) = 0$ :

Set  $A = S/I$ ,  $\mathfrak{M} = \mathfrak{N}/I$ ,  $W := H_{\mathfrak{M}}^0(A) := U/I$  and  $B := A/W = S/U$ . Then  $B$  is a hypersurface because it is Cohen–Macaulay and  $e(B) = 2$ . Thus there exists  $f \in \mathfrak{N}^2 \setminus \mathfrak{N}^3$  such that  $U = fS$ . Since  $f\mathfrak{N} \subseteq I \subseteq fS$ , we have  $I = f\mathfrak{N}$ . In particular,  $q = 3$ . Thus  $A$  has minimal multiplicity of degree 3 and  $c = 1$  by Lemma 5.1.  $\square$

**Example 5.4** (See also [4]). Let  $X_1, \dots, X_d, Y_1, \dots, Y_d, Y, Z_1, \dots, Z_c$  be indeterminates over a field  $k$ .

- (1) Put  $A = k[X_1, \dots, X_d, Y]/(X_1Y, \dots, X_dY, Y^2)$ . Then  $A$  is a Buchsbaum  $k$ -algebra with  $\dim A = d$ ,  $\text{depth } A = 0$  and  $e(A) = 1$ . Also, it has minimal multiplicity of degree 2.
- (2) Put  $A = k[X_1, \dots, X_d, Y_1, \dots, Y_d]/(X_1, \dots, X_d) \cap (Y_1, \dots, Y_d)$ . Then  $A$  is a Buchsbaum  $k$ -algebra with  $\dim A = d$ ,  $\text{depth } A = 1$  and  $e(A) = 2$ . Also, it has minimal multiplicity of degree 2.
- (3) Put  $S = k[X_1, \dots, X_d, Y_1, \dots, Y_d, Z_1, \dots, Z_c]$ . Also, if we put  $I = (X_1, \dots, X_d) \cap (Y_1, \dots, Y_d) + (Z_1, \dots, Z_c)\mathfrak{N}$  and  $A = S/I$ , then  $A$  is a Buchsbaum  $k$ -algebra with  $\dim A = d$ ,  $\text{depth } A = 0$ ,  $\text{codim } A = c$  and  $e(A) = 2$ . Also, it has minimal multiplicity of degree 2.

- (4) Put  $A = k[X_1, \dots, X_d, Y]/(X_1Y^2, \dots, X_dY^2)$ . Then  $A$  is a Buchsbaum  $k$ -algebra with  $\dim A = d$ ,  $\text{depth } A = 0$  and  $e(A) = 2$ . Also, it has minimal multiplicity of degree 3.

By a similar argument as in the proof of Proposition 5.3 one can show that any non-Cohen–Macaulay Buchsbaum  $k$ -algebra  $A$  with  $e(A) = 3$  has minimal multiplicity in our sense whenever  $q \geq 3$ ; see also Examples 5.5, 5.6. However, this is *not* true in general when  $q = 2$ ; see Example 5.6(2).

**Example 5.5** (See also Lemma 5.1). Let  $U$  be a graded ideal of  $S$  such that  $B = S/U$  is Cohen–Macaulay with  $e(B) = 3$ . If we put  $A = S/\mathfrak{M}U$ , then  $A$  is Buchsbaum such that  $A/H_{\mathfrak{M}}^0(A) \cong B$  and  $e(A) = e(B) = 3$ . Then either one of the following two cases occurs:

- (1)  $B$  has maximal embedding dimension and  $A$  has minimal multiplicity of degree 3.
- (2)  $B$  is a hypersurface and  $A$  has minimal multiplicity of degree 4.

There exists a Buchsbaum  $k$ -algebra  $A$  having minimal multiplicity with  $\text{depth } A = 0$  such that  $A/H_{\mathfrak{M}}^0(A)$  is not Cohen–Macaulay; see the next example (3).

**Example 5.6.** Let  $S = k[X, Y, Z, W]$  be a polynomial ring.

- (1) If we set  $I = (X, Y, Z, W)(XW - YZ, Y^2 - XZ, Z^2 - YW)$  and  $A = S/I$ , then  $A/H_{\mathfrak{M}}^0(A) \cong k[s^3, s^2t, st^2, t^3]$  is Cohen–Macaulay with 2-linear resolution. Thus  $A$  is Buchsbaum with minimal multiplicity of degree 3. Also, we have:

$$H_{\mathfrak{M}}^0(A) = k(-2), \quad H_{\mathfrak{M}}^1(A) = 0 \quad \text{and} \quad e(A) = 3.$$

- (2) If we set  $I = (X, Y, Z, W)(YZ, Z^2) + (Y^2 - XZ)$  and  $A = S/I$ , then  $A/H_{\mathfrak{M}}^0(A) \cong k[X, Y, Z, W]/(Y^2 - XZ, YZ, Z^2) \cong \text{gr}_{\mathfrak{m}}(k[[t^3, t^4, t^5]])[W]$  is Cohen–Macaulay with  $e(B) = 3$ . Thus  $A$  is Buchsbaum with  $\text{indeg } A (= \text{indeg } B) = 2$  and  $e(A) = 3$ . But  $A$  does *not* have minimal multiplicity and

$$H_{\mathfrak{M}}^0(A) = k(-2)^{\oplus 2}, \quad H_{\mathfrak{M}}^1(A) = 0 \quad \text{and} \quad e(A) = 3.$$

- (3) If we set  $I = ((X, Y, Z, W)(XW - YZ), Z^3 - YW^2, Y^3 - X^2Z, XZ^2 - Y^2W)$  and  $A = S/I$ , then  $A$  is Buchsbaum with minimal multiplicity of degree 3 since

$$H_{\mathfrak{M}}^0(A) = k(-2), \quad H_{\mathfrak{M}}^1(A) = k(-1) \quad \text{and} \quad e(A) = 4.$$

But  $A/H_{\mathfrak{M}}^0(A) \cong k[s^4, s^3t, st^3, t^4]$  is *not* Cohen–Macaulay.

**5.2. The case of Stanley–Reisner rings.** One can find abundant examples of Buchsbaum reduced  $k$ -algebras with minimal multiplicity in the class of Stanley–Reisner rings. In fact, in [16] Terai and the second author [16] gave another definition of Buchsbaum  $k$ -algebras with minimal multiplicity for Stanley–Reisner rings and have characterized those rings. This paper was inspired their work.

Now let us briefly recall the theory of Stanley–Reisner rings. Let  $\Delta$  be a *simplicial complex* on the vertex set  $V = [v] := \{1, \dots, v\}$ , that is,  $\Delta$  is a collection of subsets of  $V$  such that (a)  $F \subseteq G$ ,  $G \in \Delta \implies F \in \Delta$  and (b)  $\{i\} \in \Delta$  for  $i \in V$ . The dimension of  $F \in \Delta$  ( $F$  is said to be a *face* of  $\Delta$ ) is defined by  $\#(F) - 1$ . Set  $\dim \Delta = \max\{\dim F : F \in \Delta\}$ . A complex is called *pure* if all facets (maximal faces) have the same dimension. Put  $\text{link}_{\Delta} G = \{F \in \Delta : F \cup G \in \Delta, F \cap G \notin \Delta\}$ , the *link* of a face  $G$  in  $\Delta$ .

Set  $I_{\Delta} = (X_{i_1} \cdots X_{i_p} : 1 \leq i_1 < \cdots < i_p \leq v, \{i_1, \dots, i_p\} \notin \Delta)S$ . Then  $k[\Delta] = S/I_{\Delta}$  is called the *Stanley–Reisner ring* of  $\Delta$  over  $k$ . The ring  $k[\Delta]$  is a homogeneous reduced  $k$ -algebra with  $d := \dim k[\Delta] = \dim \Delta + 1$ . Moreover,  $q = \text{indeg } k[\Delta] \leq d + 1$  always holds, and  $\Delta$  is  $(d - 1)$ -skeleton of  $(v - 1)$ -simplices of  $2^{[v]}$  whenever  $q = d + 1$ . Put

$f_i = \#\{F \in \Delta : \dim F = i\}$  for all  $i \in \mathbb{Z}$ . Then the Hilbert series of  $k[\Delta]$  can be written as in the following form:

$$F(k[\Delta], t) = \sum_{i=-1}^{d-1} \frac{f_i t^{i+1}}{(1-t)^{i+1}} = \frac{h_0 + h_1 t + \cdots + h_d t^d}{(1-t)^d}.$$

In particular,  $e(k[\Delta])$  is equal to  $f_{d-1}(\Delta)$ , the number of facets  $F$  with  $\dim F = d-1$ .

In the following, let  $\Delta$  be a simplicial complex on  $V = [v]$  with  $\dim \Delta = d-1$ . For Buchsbaumness of Stanley–Reisner rings, the following fact is well-known. Note that  $k[\Delta]$  is Buchsbaum if and only if it has (F.L.C.), that is, every local cohomology modules  $H_{\mathfrak{M}}^i(k[\Delta])$  has finite length for  $i \neq d$ .

**Fact 5.7.** *The following conditions are equivalent:*

- (1)  $k[\Delta]$  is Buchsbaum.
- (2)  $\Delta$  is pure and  $k[\text{link}_{\Delta}\{i\}]$  is Cohen–Macaulay (of dimension  $d-1$ ) for each  $i \in V$ .
- (3)  $H_{\mathfrak{M}}^i(A) = [H_{\mathfrak{M}}^i(A)]_0 (\cong \tilde{H}_{i-1}(\Delta; k))$  for all  $i < d$ .

In the class of Buchsbaum Stanley–Reisner rings, there exists a criterion for  $k[\Delta]$  to have linear resolution in terms of  $h$ -vectors as follows:

**Theorem 5.8** ([16, Theorem 1.3]). *Suppose that  $A = k[\Delta]$  is Buchsbaum and  $2 \leq q \leq d$ . Put  $h = \dim_k H_{\mathfrak{M}}^{q-1}(A)$ . Then the following conditions are equivalent:*

- (1)  $A$  has  $q$ -linear resolution.
- (2) The  $h$ -vector  $h(\Delta) = (h_0, h_1, \dots, h_{q-1}, h_q, h_{q+1}, \dots, h_d)$  of  $\Delta$  is

$$\left(1, c, \dots, \binom{c+q-2}{q-1}, -\binom{d}{q}h, \binom{d}{q+1}h, \dots, (-1)^{d-q+1} \binom{d}{d}h\right)$$

- (3) The following equalities hold:

$$e(A) = \binom{c+q-1}{q-1} - h \binom{d-1}{q-1} \quad \text{and} \quad I(A) = h \binom{d-1}{q-1}.$$

When this is the case,  $H_{\mathfrak{M}}^i(A) = 0$  ( $i \neq q-1, d$ ) and the following inequalities hold:

$$0 \leq h \leq \frac{c(c+1) \cdots (c+q-2)}{d(d-1) \cdots (d-q+2)} =: h_{c,d,q}.$$

From this point of view, one can regard a *Buchsbaum simplicial complex with minimal multiplicity* as a Buchsbaum complex with linear resolution and “maximal homology”.

**Theorem 5.9** ([16, Theorem 2.3]). *Suppose that  $A = k[\Delta]$  is Buchsbaum. Then*

$$e(A) \geq \frac{c+d}{d} \binom{c+q-2}{q-2}$$

*holds, and the following statements are equivalent:*

- (1)  $\Delta$  is a Buchsbaum complex with minimal multiplicity of degree  $q$ , that is,  $e(A) = \frac{c+d}{d} \binom{c+q-2}{q-2}$ .
- (2)  $A$  has minimal multiplicity of degree  $q$  in our sense.
- (3)  $A$  has  $q$ -linear resolution with  $l_A(H_{\mathfrak{M}}^{q-1}(A)) = h_{c,d,q}$ .
- (4)  $k[\text{link}_{\Delta}\{i\}]$  has  $(q-1)$ -linear resolution for all  $i \in V$ .
- (5) The  $h$ -vector of  $A$  can be written as in the form of Theorem 5.8(2) and  $h = h_{c,d,q}$ .
- (6)  $k[\Delta^*]$  has pure and almost linear resolution and  $a(k[\Delta^*]) = 0$ , where  $\Delta^* = \{F \in \Delta : V \setminus F \notin \Delta\}$  denotes the Alexander dual complex of  $\Delta$ :

$$0 \rightarrow S(-(c+d))\beta_q^* \rightarrow S(-(c+q-2))\beta_{q-1}^* \rightarrow \cdots \rightarrow S(-c)\beta_1^* \rightarrow S \rightarrow k[\Delta^*] \rightarrow 0,$$

where  $S = k[X_1, \dots, X_v]$  and  $\beta_q^* = h_{c,d,q}$ .

Let us pick up some examples of Buchsbaum complexes with minimal multiplicity from [16].

**Example 5.10.**  $\Delta$  is a Buchsbaum complex with minimal multiplicity of degree 2 if and only if it is a finite disjoint union of  $(d-1)$ -simplices.

When this is the case,  $k[\Delta]$  is isomorphic to

$$k[X_{ij} : 1 \leq i \leq d, 1 \leq j \leq e] / (X_{pr}X_{qs} : 1 \leq p, q \leq d, 1 \leq r \neq s \leq e),$$


where  $e = e(k[\Delta])$ , the number of connected components of  $\Delta$ .

Now let us recall a cyclic polytope. The algebraic curve  $M \subseteq \mathbb{R}^f$  parametrically by  $x(t) = (t, t^2, \dots, t^f)$  is called the moment curve. Let  $n \geq f+1$  be an integer. A *cyclic polytope* with  $n$  vertices, denoted by  $C(n, f)$ , is the convex hull of any  $n$  distinct points on  $M$ . Note that  $C(n, f)$  is a simplicial  $f$ -polytope. Next example provides us  $d$ -dimensional Buchsbaum reduced  $k$ -algebras of degree  $q$  for given integers  $2 \leq q \leq d$ .

**Example 5.11.** Let  $q, d$  be given integers with  $2 \leq q \leq d$ . Put  $n = 2d - q + 2$  and  $f = 2(d - q + 1)$ . Let  $\Delta$  be the Alexander dual of the boundary complex  $\Gamma$  of a cyclic polytope  $C(n, f)$ . Then since  $k[\Gamma]$  satisfies the condition (6) in Theorem 5.9, we obtain that  $k[\Delta]$  is a  $d$ -dimensional Buchsbaum Stanley–Reisner ring with minimal multiplicity of degree  $q$  with  $h = h_{c,d,q} = 1$ .

For a given integer  $n \geq 5$  such that  $n \equiv 0, 2 \pmod{3}$ , there exists a 2-dimensional Buchsbaum complex on  $[n]$  with minimal multiplicity of degree 3; see [9, 16, 17].

**Example 5.12.** Let  $\Delta$  be a simplicial complex on  $V = [5]$  spanned by  $\{1, 2, 3\}$ ,  $\{1, 3, 4\}$ ,  $\{1, 4, 5\}$ ,  $\{2, 3, 5\}$  and  $\{2, 4, 5\}$  as follows:



Then

$$\begin{aligned} k[\Delta] &= k[X_1, X_2, X_3, X_4, X_5] / (X_1X_2X_4, X_1X_2X_5, X_1X_3X_5, X_2X_3X_4, X_3X_4X_5) \\ &= k[X_1, X_2, X_3, X_4, X_5] / (X_1, X_4) \cap (X_4, X_5) \cap (X_2, X_5) \cap (X_2, X_3) \cap (X_1, X_3) \end{aligned}$$

is a 3-dimensional Buchsbaum rings with minimal multiplicity of degree 3.

In general, the following problem remains open when  $d \geq 4$ . When  $q = d$ , it suffices to show the existence of  $(d-1)$ -dimensional Buchsbaum complexes with minimal multiplicity of degree  $q$  with  $\text{codim } k[\Delta] = c$ ; see [17].

**Problem 5.13.** Let  $c, d, q, h$  be integers with  $c \geq 2$ ,  $2 \leq q \leq d$  and  $0 \leq h \leq h_{c,d,q}$ . Construct  $(d-1)$ -dimensional Buchsbaum complexes  $\Delta$  with  $q$ -linear resolution such that  $\text{codim } k[\Delta] = c$  and  $\dim_k H_{\mathfrak{M}}^{q-1}(k[\Delta]) = h$ .

**5.3. The case of integral domains.** There are many examples of homogeneous Buchsbaum integral domains having linear resolutions; see e.g., [1, 3, 7]. However we have no examples of Buchsbaum homogeneous integral domain over an algebraically closed field  $k$  with minimal multiplicity of degree  $q \geq 2$ . Indeed, our examples described before are not integral domains.

**Discussion 5.14** (See [3]). Let  $d \geq 2$ ,  $h^i$  ( $1 \leq i \leq d-1$ ),  $s \geq 0$  be integers. Let  $k$  an algebraically closed field. Then there exist  $d$ -dimensional Buchsbaum homogeneous  $k$ -algebra  $A$  which are integral domains having  $q$ -linear resolutions with  $\text{codim } A = 2$  and such

that  $\dim_k H_{\mathfrak{M}}^i(A) = h^i$  for all  $i = 1, \dots, d-1$ , where

$$q = \sum_{i=1}^{d-1} \left\{ \sum_{j=1}^{d+2-i} (-1)^j \cdot (j-2) \cdot \binom{d+2}{i+j} \right\} h^i + s - 3.$$

But this number  $q$  is too big to satisfy the condition (4) in Theorem 4.1.

The following proposition is maybe well-known for experts.

**Proposition 5.15.** *Let  $A$  be a Buchsbaum homogeneous  $k$ -algebra with minimal multiplicity of degree at most 2. Suppose that  $k$  is algebraically closed and that  $A$  is an integral domain. Then  $A$  is isomorphic to a polynomial ring.*

*Proof.* By Abhyankar's result mentioned as in the introduction, we have  $\text{embdim}(A) \leq e(A) + \dim A - 1$ . On the other hand,  $\text{embdim}(A) = e(A) + \dim A - 1 + I(A)$  since  $A$  has maximal embedding dimension. Hence  $I(A) = 0$ , that is,  $A$  is Cohen–Macaulay. Thus  $A$  is a polynomial ring by Corollary 4.4.  $\square$

Based on the above observation we pose the following conjecture:

**Conjecture 5.16.** *There is no Buchsbaum homogeneous integral domain over an algebraically closed field with minimal multiplicity of degree  $q \geq 2$ .*

In the above conjecture we cannot remove the assumption that  $k$  is algebraically closed.

**Example 5.17** (See [4]). Let  $K/k$  be an extension of fields of degree 2, and set  $B = K[X_1, \dots, X_n]$ , a polynomial ring. Then  $A = \{f \in B : f(0, \dots, 0) \in k\}$  is a homogeneous Buchsbaum  $k$ -algebra with minimal multiplicity of degree 2 since  $e(A) = 2$ .

## REFERENCES

- [1] M. Amasaki, Applications of the generalized Weierstrass preparation theorem to the study of homogeneous ideals, *Trans. Amer. Math. Soc.* **317** (1990), 1–43.
- [2] S. S. Abhyankar, *Local rings of high embedding dimension*, *Amer. J. Math.* **89** (1967), 1073–1077.
- [3] D. Eisenbud and S. Goto, *Linear free resolutions and minimal multiplicity*, *J. Algebra* **88** (1984), 89–133.
- [4] S. Goto, *Buchsbaum rings with multiplicity 2*, *J. Algebra* **74** (1982), 494–508.
- [5] S. Goto, *Buchsbaum rings of maximal embedding dimension*, *J. Algebra* **76** (1982), 383–399.
- [6] S. Goto, *On the associated graded rings of parameter ideals in Buchsbaum rings*, *J. Algebra* **85** (1983), 490–534.
- [7] S. Goto, *Noetherian local rings with Buchsbaum associated graded rings*, *J. Algebra* **86** (1984), 336–384.
- [8] S. Goto and K. Yoshida, *Buchsbaum local rings with minimal multiplicity*, in preparation.
- [9] K. Hanano, *Construction of two-dimensional Buchsbaum simplicial complexes*, *Europ. J. Combinatorics* **22** (2001), 171–178.
- [10] L. T. Hoa and C. Miyazaki, *Bounds on Castelnuovo–Mumford regularity for generalized Cohen–Macaulay graded rings*, *Math. Ann.* **301** (1995), 587–598.
- [11] M. Herrmann and S. Ikeda, *Three notes on the order of ideals defining hypersurfaces*, *J. Algebra* **132** (1990), 123–129.
- [12] Y. Kamoi and W. Vogel, *On Buchsbaum Algebras*, *J. Pure Appl. Algebra* **122** (1997), 251–263.
- [13] J. D. Sally, *On the associated graded rings of a local Cohen–Macaulay ring*, *J. Math. Kyoto Univ.* **17** (1977), 19–21.
- [14] J. D. Sally, *Cohen–Macaulay local rings of maximal embedding dimension*, *J. Algebra* **56** (1979), 168–183.
- [15] J. Stückrad and W. Vogel, *Buchsbaum Rings and Applications*, Springer–Verlag, Berlin, Heidelberg, New York, 1986.
- [16] N. Terai and K. Yoshida, *Buchsbaum Stanley–Reisner rings with minimal multiplicity*, *Proc. Amer. Math. Soc.* **134** (2006), 55–65.
- [17] N. Terai and K. Yoshida, *Buchsbaum Stanley–Reisner rings and Cohen–Macaulay covers*, (to appear in *Comm. in Algebra*).
- [18] K. Yamagishi, *Idealizations of maximal Buchsbaum modules over a Buchsbaum ring*, *Math. Proc. Camb. Phil. Soc.* **104** (1988), 451–478.

DEPARTMENT OF MATHEMATICS, SCHOOL OF SCIENCE AND TECHNOLOGY, MEIJI UNIVERSITY, 214-8571,  
JAPAN

*E-mail address:* goto@math.meiji.ac.jp

GRADUATE SCHOOL OF MATHEMATICS, NAGOYA UNIVERSITY, NAGOYA 464-8602, JAPAN

*E-mail address:* yoshida@math.nagoya-u.ac.jp